






PAGER Version 5

Created: 1 week, 4 days after earthquake

M 5.5, 53 km N of Miyako, Japan

Origin Time: 2020-12-12 07:18:54 UTC (Sat 16:18:54 local) Location: 40.1211° N 142.0549° E Depth: 44.0 km

Estimated Fatalities 10,000 1,000



Estimated Population Exposed to Earthquake Shaking

ESTIMATED POPULATION EXPOSURE (k=x1000)		_*	5,453k	331k	0	0	0	0	0	0
ESTIMATED MODIFIED MERCALLI INTENSITY		I	11-111	IV	V	VI	VII	VIII	IX	X+
PERCEIVED SHAKING		Not felt	Weak	Light	Moderate	Strong	Very Strong	Severe	Violent	Extreme
POTENTIAL DAMAGE	Resistant Structures	None	None	None	V. Light	Light	Moderate	Mod./Heavy	Heavy	V. Heavy
	Vulnerable Structures	None	None	None	Light	Moderate	Mod./Heavy	Heavy	V. Heavy	V. Heavy

^{*}Estimated exposure only includes population within the map area.

Population Exposure

population per 1 sq. km from Landscan

5000 10000 142.2°E 143.6°E Ĥachinohe Morioka 1iyako Mizusawa Ichinose

PAGER content is automatically generated, and only considers losses due to structural damage. Limitations of input data, shaking estimates, and loss models may add uncertainty. https://earthquake.usgs.gov/earthquakes/eventpage/us7000cqhy#pager

Structures

Overall, the population in this region resides in structures that are resistant to earthquake shaking, though vulnerable structures exist. The predominant vulnerable building types are heavy wood frame and reinforced/confined masonry construction.

Historical Earthquakes

Date	Dist.	Mag.	Max	Shaking
(UTC)	(km)		MMI(#)	Deaths
1994-12-28	126	7.7	VII(130k)	3
1983-05-26	254	7.7	VII(174k)	104
1993-07-12	387	7.7	VIII(4k)	200

Recent earthquakes in this area have caused secondary hazards such as landslides and fires that might have contributed to losses.

Selected City Exposure

from GeoNames.org MMI City Population IV Uchimaru <1kI۷ **Hachinohe** 239k IV Misawa 43k Ш Miyako 52k Ш **Furudate** <1kШ Inuotose <1kШ Morioka 295k

Ш **Aomorishi** 298k Ш Akita 326k Ш Akita 320k Ш Hakodate 276k

bold cities appear on map.

(k = x1000)